

ORIGINAL ARTICLE

International spinal cord injury cardiovascular function basic data set

A Krassioukov¹, MS Alexander², A-K Karlsson³, W Donovan⁴, CJ Mathias^{5,6} and F Biering-Sørensen⁷

¹International Collaboration On Repair Discoveries (ICORD), Department of Medicine, University of British Columbia and Vancouver Coastal Health, Vancouver, British Columbia, Canada; ²Physical Medicine and Rehabilitation, University of Alabama, Birmingham, AL, USA; ³Institute of Neuroscience and Physiology, Sahlgrenska University Hospital, Gothenburg, Sweden; ⁴Physical Medicine and Rehabilitation, University of Texas Health Science Center, Baylor College of Medicine, Houston, TX, USA; ⁵Neurovascular Medicine Unit, Faculty of Medicine, Imperial College London, London, UK; ⁶Autonomic Unit, National Hospital for Neurology and Neurosurgery, Institute of Neurology, Queen Square, University College London, London, UK and ⁷Clinic for Spinal Cord Injuries, Clinic for Spinal Cord Injuries, NeuroScience Centre, Rigshospitalet, and University of Copenhagen, Copenhagen, Denmark

Objective: To create an International Spinal Cord Injury (SCI) Cardiovascular Function Basic Data Set within the framework of the International SCI Data Sets.

Setting: An international working group.

Methods: The draft of the data set was developed by a working group comprising members appointed by the American Spinal Injury Association (ASIA), the International Spinal Cord Society (ISCoS) and a representative of the executive committee of the International SCI Standards and Data Sets. The final version of the data set was developed after review by members of the executive committee of the International SCI Standards and Data Sets, the ISCoS scientific committee, ASIA board, relevant and interested international organizations and societies, individual persons with specific interest and the ISCoS Council. To make the data set uniform, each variable and each response category within each variable have been specifically defined in a way that is designed to promote the collection and reporting of comparable minimal data.

Results: The variables included in the International SCI Cardiovascular Function Basic Data Set include the following items: date of data collection, cardiovascular history before the spinal cord lesion, events related to cardiovascular function after the spinal cord lesion, cardiovascular function after the spinal cord lesion, medications affecting cardiovascular function on the day of examination; and objective measures of cardiovascular functions, including time of examination, position of examination, pulse and blood pressure. The complete instructions for data collection and the data sheet itself are freely available on the websites of both ISCoS (<http://www.iscos.org.uk>) and ASIA (<http://www.asia-spinalinjury.org>).

Spinal Cord (2010) 48, 586–590; doi:10.1038/sc.2009.190; published online 26 January 2010

Keywords: spinal cord injury; international data set; cardiovascular function

Introduction

Cardiovascular abnormalities have been well documented in humans after spinal cord lesions.^{1,2} The recognition and management of these cardiovascular dysfunctions after spinal cord injury (SCI) represent challenging clinical issues. Moreover, cardiovascular disorders in the acute and chronic stages of SCI are among the most common causes of death in individuals with SCI.^{3–5} Unfortunately, little attention has been paid to the documentation of these dysfunctions in individuals with spinal cord lesions.⁶

In accordance with the aims of the International SCI Data Sets,⁷ the aim of the Cardiovascular Function Basic Data Set

for SCI is to standardize the collection and reporting of a minimal amount of information on cardiovascular function in daily practice. Furthermore, the International SCI Cardiovascular Function Basic Data Set makes it possible to evaluate and compare results from various published studies on cardiovascular function after SCI.

The International SCI Cardiovascular Function Basic Data Set is applicable to adults with traumatic or non-traumatic supraconal, conal or cauda equina lesions. To ensure that data are standardized, each variable and each response category within variables have been specifically defined. The Cardiovascular Function Basic SCI Data Set will be used in connection with the background information within the International SCI Core Data Set.⁸ The SCI Core Data Set documents the level, completeness and time post-spinal cord lesion, which have an important role in cardiovascular outcomes after injury.

Correspondence: Dr A Krassioukov, International Collaboration On Repair Discoveries (ICORD), UBC, 818 West 10th Avenue, Vancouver, British Columbia, Canada V5Z 1M9.

E-mail: krassioukov@icord.org

Received 15 September 2009; revised and accepted 1 December 2009; published online 26 January 2010

Materials and methods

The first draft of the International SCI Cardiovascular Function Basic Data Set was made by a working group consisting of members appointed by the American Spinal Injury Association (ASIA) and the International Spinal Cord Society (ISCOs), together with a representative of the Executive Committee of the International SCI Standards and Data Sets. The developmental process for the International Cardiovascular Function Basic SCI Data Set followed the steps given below:

- (1) The working group of the International SCI Cardiovascular Function Basic Data Set finalized the first draft during a 2 day meeting in Copenhagen in March 2007. This was further elaborated by frequent e-mail contacts between the group members.
- (2) The data set has been reviewed by members of the executive committee of the International SCI Standards and Data Sets.
- (3) The comments from the committee members were discussed in the working group and appropriate responses were made to the data set.
- (4) Members of the ISCOs scientific committee and ASIA board were also asked to review the data set.
- (5) The comments from the committee and board members were discussed in the working group and a response was made, and further adjustments of the data set were performed.
- (6) Relevant and interested scientific and professional international organizations and societies (International Society of Physical Medicine and Rehabilitation, American Paraplegia Society; and others) and individuals who were interested were also invited to review the data set. In addition, the data set was posted on the ISCOs and ASIA websites for over 2 months to allow comments and suggestions.
- (7) The comments were discussed and responded to by the working group and wherein appropriate, adjustments to the data set were made.

Results

The International SCI Cardiovascular Function Basic Data Set is structured according to established protocol for the International SCI Data Sets.⁷ The complete data set form is included in the Appendix. The complete data syllabus, data sheet and training cases will be available at the respective websites of ISCOs (www.iscos.org.uk) and ASIA (www.asia-spinalinjury.org).

Date of data collection

As the collection of data on cardiovascular functions may be conducted at any time after SCI, the date of data collection is imperative to compute the length of time since the spinal cord lesion occurred and to identify data collected in relation to other data collected on the same individual at various time points.

Cardiovascular history before spinal cord lesion

This variable will document the history of cardiovascular function present before the spinal cord lesion and should be

collected only once. Cardiovascular abnormalities present in an individual before SCI is a major concern as additional deterioration of cardiovascular function could occur as a result of SCI.

Abnormal heart rates (HRs) and rhythms are commonly present after a spinal cord lesion.^{9,10} Presence of a cardiac pacemaker, previous surgeries (for example, ablation of ectopic foci) or other conditions (for example, pre-existing atrial fibrillation, myocardial infarction or congestive heart failure) could influence these parameters.¹¹

Pre-injury the level of arterial blood pressure (BP) could be either low because of hypotension or elevated because of hypertension. Altered autonomic control after spinal cord lesion could further exacerbate preexisting instability of BP. Hypotension (systolic arterial BP <90 mm Hg) is common in acute and chronic spinal cord lesions.^{12–14} Addition, intermittent hypertension can be associated with noxious or non-noxious stimuli and resultant autonomic dysreflexia.² Preexisting abnormalities of BP can influence the cardiovascular functions after spinal cord lesion.¹⁵

Hypertension. (arterial BP >140/90 mm Hg),¹⁵ preexisting elevation in arterial BP, not associated with episodes of autonomic dysreflexia.

Orthostatic hypotension is symptomatic or asymptomatic decrease in BP usually exceeding 20 mm Hg systolic or 10 mm Hg diastolic on moving from the supine to an upright position.¹⁶

Deep vein thrombosis of the legs, pelvis or arms, because of coagulopathy, stasis or endothelial injury are common in individuals with spinal cord lesion.¹⁷

Preexisting *neuropathies* (for example, diabetic or alcoholic neuropathy) and *other conditions* associated with autonomic dysfunctions (for example, Parkinson's disease, multiple sclerosis and traumatic brain injury) can affect altered cardiovascular functions post-spinal cord lesion, and should therefore be included.^{16,18–20}

Events related to cardiovascular function after spinal cord lesion

Any events related to cardiovascular functions after spinal cord lesion should be documented. These time-limited cardiovascular events with long-term sequelae should be dated to enable computation of time since injury and to identify the relationship between data points.

Cardiovascular function history after spinal cord lesion within the last 3 months *Cardiac conditions* with subjective symptoms related to the heart that occur post-spinal cord lesion should be documented (for example, abnormal HRs/rhythm, angina, palpitation, and so on).

Orthostatic hypotension similarly as above.¹⁶

Dependent oedema is a clinically detectable increase in extracellular fluid volume localized in a dependent area, such as a limb, characterized by swelling or pitting.¹⁵

Hypertension. The diagnosis of *hypertension* (arterial BP >140/90 mm Hg) in individuals with SCI should be considered after careful monitoring and documentation of arterial BP and exclusion of elevation of BP because of episodes of autonomic dysreflexia.

Autonomic dysreflexia is a constellation of signs and/or symptoms in persons with SCI above T5-6 in response to noxious or non-noxious stimuli below the level of injury. Autonomic dysreflexia is characterized by an increase in systolic BP (>20 mmHg above baseline), and may include one of the following symptoms: headache, flushing and sweating above the level of the lesion, vasoconstriction below the level of the lesion, and dysrhythmia.^{1,2,21-23} This syndrome may or may not be symptomatic and may occur at any time after SCI.^{24,25}

Medication affecting cardiovascular function on the day of examination. All medications affecting cardiovascular functions (for example, HR, BP), which are presently taken by the individual should be documented.

The final part of the International SCI Cardiovascular Function Basic Data Set contains true cardiovascular data that should be collected on the day of examination.

Time performed. Cardiovascular parameters are affected by circadian rhythms, therefore, the exact time of evaluation should be reported to appreciate this variability.^{26,27}

Position during testing. Cardiovascular parameters similarly are affected by the position of the individual during the testing. For example, BP could decrease because of sitting or standing position.^{14,28}

Devices in use during the testing. The wearing of compression devices (abdominal binder or pressure stockings) could affect cardiovascular parameters during the examination,^{29,30} therefore, their use during the examination should be documented.

Pulse. HR is a standard cardiovascular parameter documented during evaluations. The time after injury, as well as level and completeness of spinal cord lesion are crucial factors that affect HR. Individuals with injuries at T6 and below have preserved sympathetic and parasympathetic control to the heart and do not show HR abnormalities related to spinal cord lesion.³¹ Both abnormal HR and rhythm are commonly observed in individuals with cervical and high thoracic spinal cord lesions.^{10,32,33}

Blood pressure. Low resting arterial BP is common in individuals with spinal cord lesions at T6 and above.^{34,35} Furthermore, these individuals are also prone to abnormal arterial BP fluctuation because of orthostatic instability or episodes of autonomic dysreflexia. Individuals with lesions at the lower thoracic spinal cord or conus medullaris usually show normal arterial BP because of preserved sympathetic control to the heart and splanchnic circulation.^{1,2,36} Resting arterial BP and HR should be obtained after voiding and 5 min of rest. Both physical activity and full urinary bladder could affect cardiovascular parameters.³⁷

Discussion

The International SCI Cardiovascular Function Basic Data Set incorporates the data that shall be observed in conjunction with

data in the International SCI Core Data Set⁸ and other relevant autonomic functions data sets (Pulmonary, Skin/Thermoregulation/Sudomotor and Endocrine/Metabolic Data Set) that are presently in development. The International SCI Core Data Set includes, among other things, information on the date of birth and injury, gender, the cause of spinal cord lesion and neurological status, whereas the International SCI Cardiovascular Function Basic Data set includes variables on date of data collection, cardiovascular history before the spinal cord lesion, events related to cardiovascular functions after the spinal cord lesion, cardiovascular functions after the spinal cord lesion, medications affecting cardiovascular functions on the day of examination; and objective measures of cardiovascular functions (time of examination, position of examination, BP and HR).

It is understood that the arterial BP and HR are very sensitive to various intrinsic and extrinsic factors including preexisting cardiovascular diseases, medications, time of day and position in which the measures were taken.^{29,38,39} Accordingly, the International SCI Cardiovascular Function Basic Data Set includes data on numerous variables that could influence interpretation of the cardiovascular parameters collected in individual with SCI on the day of examination.

Abnormal cardiovascular control should be expected in individuals with spinal cord lesion of both traumatic¹ or non-traumatic etiology.^{40,41} Although, the most prominent cardiovascular dysfunctions are observed in individuals with cervical and upper thoracic lesions all lesions to the spinal cord including conus medullaris and cauda equina are included in this context. Even individuals with lower levels of spinal cord trauma could experience cardiovascular dysfunctions in the early stages after the SCI.¹²

It is extremely important that data be collected in a uniform manner. Furthermore, the use of a standard format is essential for combining and comparing the data from multiple sites. For these reasons, each variable and each response category within each variable have been specifically defined in a way that is designed to promote the collection and reporting of comparable minimal data. It is expected that more detailed information on cardiovascular parameters will be provided when using the International SCI Cardiovascular Function Extended Data Set. This information will be probably too extensive in the typical clinical setting and is mainly intended for clinical studies.

Although, the International SCI Cardiovascular Function Basic Data Set has been revised by the international community it is expected that this data set will require periodic revisions and updates. Ideas for improvement of the data set are welcome and should be forwarded to the corresponding author.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

Coloplast A/S, Denmark has supported the work for this data set with an unconditional grant. We are thankful for comments and suggestions received from Dr C Ho, Dr S Charlifue, Dr E Inge, Dr L Vogel and Dr G Müller. This

International SCI Cardiovascular Function Data Set will be announced at the websites of ISCoS (<http://www.iscos.org.uk>) and ASIA (<http://www.asia-spinalinjury.org>).

References

- Krassioukov A, Claydon VE. The clinical problems in cardiovascular control following spinal cord injury: an overview. *Prog Brain Res* 2006; **152**: 223–229.
- Mathias CJ, Frankel HL. Autonomic disturbances in spinal cord lesions. In: Bannister R, Mathias CJ (eds). *Autonomic Failure, a Textbook of Clinical Disorders of the Autonomic Nervous System*. Oxford University Press: New York, 2002, pp 839–881.
- DeVivo MJ, Krause JS, Lammertse DP. Recent trends in mortality and causes of death among persons with spinal cord injury. *Arch Phys Med Rehabil* 1999; **80**: 1411–1419.
- Garshick E, Kelley A, Cohen SA, Garrison A, Tun CG, Gagnon D et al. A prospective assessment of mortality in chronic spinal cord injury. *Spinal Cord* 2005; **43**: 408–416.
- Bauman WA, Spungen AM. Coronary heart disease in individuals with spinal cord injury: assessment of risk factors. *Spinal Cord* 2008; **46**: 466–476.
- Krassioukov AV, Karlsson AK, Wecht JM, Wuermser LA, Mathias C, Marino RJ. Assessment of autonomic dysfunction following spinal cord injury: rationale for additions to the international standards for neurological assessment. *J Rehabil Res Dev* 2007; **44**: 103–112.
- Biering-Sorensen F, Charlifue S, DeVivo M, Noonan V, Post M, Stripling T et al. International spinal cord injury data sets. *Spinal Cord* 2006; **44**: 530–534.
- DeVivo M, Biering-Sorensen F, Charlifue S, Noonan V, Post M, Stripling T et al. International spinal cord injury core data set. *Spinal Cord* 2006; **44**: 535–540.
- Gilgoff IS, Ward SL, Hohn AR. Cardiac pacemaker in high spinal cord injury. *Arch Phys Med Rehabil* 1991; **72**: 601–603.
- Franga DL, Hawkins ML, Medeiros RS, Adewumi D. Recurrent asystole resulting from high cervical spinal cord injuries. *Am Surg* 2006; **72**: 525–529.
- Kalahasty G, Ellenbogen K. The role of pacemakers in the management of patients with atrial fibrillation. *Med Clin North Am* 2008; **92**: 161–178 xi-xii.
- Sidorov EV, Townson AF, Dvorak MF, Kwon BK, Steeves J, Krassioukov A. Orthostatic hypotension in the first month following acute spinal cord injury. *Spinal Cord* 2008; **46**: 65–69.
- Illman A, Stiller K, Williams M. The prevalence of orthostatic hypotension during physiotherapy treatment in patients with an acute spinal cord injury. *Spinal Cord* 2000; **38**: 741–747.
- Claydon VE, Krassioukov AV. Orthostatic hypotension and autonomic pathways after spinal cord injury. *J Neurotrauma* 2006; **23**: 1713–1725.
- Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN et al. Recommendations for blood pressure measurement in humans and experimental animals: part 1: blood pressure measurement in humans: a statement for professionals from the subcommittee of professional and public education of the American heart association council on high blood pressure research. *Circula* 2005; **111**: 697–716.
- Consensus statement on the definition of orthostatic hypotension, pure autonomic failure, and multiple system atrophy. The consensus committee of the American autonomic society and the American academy of neurology. *Neurol* 1996; **46**: 1470.
- Prevention of thromboembolism in spinal cord injury. Consortium for spinal cord medicine. *J Spinal Cord Med* 1997; **20**: 259–283.
- Kodounis A, Stamboulis E, Constantinidis TS, Liolios A. Measurement of autonomic dysregulation in multiple sclerosis. *Acta Neurol Scand* 2005; **112**: 403–408.
- Merkelbach S, Haensch CA, Hemmer B, Koehler J, Konig NH, Ziemssen T. Multiple sclerosis and the autonomic nervous system. *J Neurol* 2006; **253**(Suppl 1): I21–I25.
- Mathias CJ. Disorders affecting autonomic function in Parkinsonian patients. In: Battistin L, Scarlato G, Caraceni T, Ruggieri S (eds). *Parkinson's Disease*. Lippincott-Raven: Philadelphia, 1996, pp 383–391.
- Krassioukov AV, Furlan JC, Fehlings MG. Autonomic dysreflexia in acute spinal cord injury: an under-recognized clinical entity. *J Neurotrauma* 2003; **20**: 707–716.
- Gao SA, Ambring A, Lambert G, Karlsson AK. Autonomic control of the heart and renal vascular bed during autonomic dysreflexia in high spinal cord injury. *Clin Auton Res* 2002; **12**: 457–464.
- Karlsson AK, Friberg P, Lonnroth P, Sullivan L, Elam M. Regional sympathetic function in high spinal cord injury during mental stress and autonomic dysreflexia. *Brain* 1998; **121**: 1711–1719.
- Kirshblum SC, House JG, O'Connor KC. Silent autonomic dysreflexia during a routine bowel program in persons with traumatic spinal cord injury: a preliminary study. *Arch Phys Med Rehabil* 2002; **83**: 1774–1776.
- Linsenmeyer TA, Campagnolo DI, Chou IH. Silent autonomic dysreflexia during voiding in men with spinal cord injuries. *J Urol* 1996; **155**: 519–522.
- Davidson C, Smith D, Morgan DB. Diurnal pattern of water and electrolyte excretion and body weight in idiopathic orthostatic hypotension. The effect of three treatments. *Am J Med* 1976; **61**: 709–715.
- Munakata M, Kameyama J, Nonukawa T, Moriai N, Yoshinaga K. Circadian blood pressure rhythm in patients with higher and lower spinal cord injury: simultaneous evaluation of autonomic nervous activity and physical activity. *J Hypertens* 1997; **15**: 1745–1749.
- Harkema SJ, Ferreira CK, van den Brand RJ, Krassioukov AV. Improvements in orthostatic instability with stand locomotor training in individuals with spinal cord injury. *J Neurotrauma* 2008; **25**: 1467–1475.
- Krassioukov AV, Harkema SJ. Effect of harness application and postural changes on cardiovascular parameters of individuals with spinal cord injury. *Spinal Cord* 2006; **44**: 780–786.
- Hopman MT, Monroe M, Dueck C, Phillips WT, Skinner JS. Blood redistribution and circulatory responses to submaximal arm exercise in persons with spinal cord injury. *Scand J Rehabil Med* 1998; **30**: 167–174.
- Lehmann KG, Shandling AH, Yusi AU, Froelicher VF. Altered ventricular repolarization in central sympathetic dysfunction associated with spinal cord injury. *Am J Cardiol* 1989; **63**: 1498–1504.
- Claydon VE, Elliott SL, Sheel AW, Krassioukov A. Cardiovascular responses to vibrostimulation for sperm retrieval in men with spinal cord injury. *J Spinal Cord Med* 2006; **29**: 207–216.
- Silbert PL, Davis MJE. Late asystole in high cervical spinal-cord injury—case-report. *Parap* 1990; **28**: 137–140.
- Mathias CJ, Frankel HL. The cardiovascular system in tetraplegia and paraplegia. In: Frankel HL (ed). *Handbook of Clinical Neurology*. B.V.: Elsevier Science Publishers: Amsterdam, Netherlands, 1992, pp 435–456.
- Sheel AW, Krassioukov AV, Inglis JT, Elliott SL. Autonomic dysreflexia during sperm retrieval in spinal cord injury: influence of lesion level and sildenafil citrate. *J A Physiol* 2005; **99**: 53–58.
- Teasell R, Arnold AP, Krassioukov AV, Delaney GA. Cardiovascular consequences of loss of supraspinal control of the sympathetic nervous system following spinal cord injuries. *Arch Phys Med Rehabil* 2000; **81**: 506–516.
- Claydon VE, Hol AT, Eng JJ, Krassioukov AV. Cardiovascular responses and postexercise hypotension after arm cycling exercise in subjects with spinal cord injury. *Arch Phys Med Rehabil* 2006; **87**: 1106–1114.
- Wecht JM, Weir JP, Krothe AH, Spungen AM, Bauman WA. Normalization of supine blood pressure after nitric oxide synthase inhibition in persons with tetraplegia. *J Spinal Cord Med* 2007; **30**: 5–9.
- Tolbert G, Tuck ML. Ambulatory blood pressure monitoring in persons with chronic spinal cord injury. *J Spinal Cord Med* 2004; **27**: 476–480.
- Vorobeychik G, Krassioukov A, Nelson J, Spring J, Beauregard N, Bozek C. Autonomic dysfunction in multiple sclerosis patients with mild disability and spinal cord lesions: reality or myth? *Multiple Sclerosis* 2006; **12**: S36.
- Flachenecker P, Wolf A, Krauser M, Hartung HP, Reiners K. Cardiovascular autonomic dysfunction in multiple sclerosis: correlation with orthostatic intolerance. *J Neurol* 1999; **246**: 578–586.

